

## Biosynthesis and chemistry of Thyroid Hormones

### CHEMISTRY :-

Thyroid gland contain an iodized glycoprotein Thyroglobulin, which apparently constitutes the storage form of the thyroid hormones. On the hydrolysis this protein yield several iodine containing derivatives of tyrosine (mono and di-iodotyrosin) and thyronine (di-, tri, and tetra iodothyronines). Among these the tetra-iodothyronine (Thyroxine) and 3<sup>5</sup>-3<sup>6</sup>-triiodothyronine are the most important. In addition to these iodine containing compounds thyroid gland also produces a non iodinated compound known as Calciferon or Thyrocalcitonin a polypeptide which lowers serum calcium concentration.

### SYNTHESIS OF THYROID HORMONES:-

The steps involve in biosynthesis of thyroid hormones are given below.

- ① SYNTHESIS OF T<sub>4</sub>G.
- ② IODINE ACCUMULATION BY THYROID.
- ③ OXIDATION OF IODIDE.
- ④ FORMATION OF ORGANICALLY BOUND IODINE COMPOUND
- ⑤ PROTEOLYSIS OF THYROGLOBULIN.

### SYNTHESIS OF THYROGLOBULIN:-

T<sub>4</sub>G is a large globular glycoprotein constituting about 75% of the protein content of mammalian Thyroid. Each molecule consist of two identical subunit and contain carbohydrates which is about 10% by weight. There are three type of polysaccharides which are covalently linked to as per gene: Serine and Threonine residues. ~~There are 140 tyrosine residue.~~

↓ α-cherry gene

(3)

(2)

28% of these residue are iodinated. T.G. contain 123 tyrosine residue but only four to eight of these one incorporated into thyroid hormone.

Thyroglobulin is synthesized by epithelial cells thyroid follicles. The TG is secreted into central cavity of follicle and constitutes the major protein component of colloid. Thyroglobulin subunits are synthesized by polyribosomes of the rough endoplasmic reticulum (RER). They assemble and are discharged through RER to golgi apparatus. During this transfer the molecule is glycosylated by specific glycosyl transferases. The completed molecule as packed into vesicles are secreted by the apical membrane into the lumen of the follicle.

Note- Thyroid cells thus have three functions. They collect and transport iodine, they synthesize TG and secrete it into the colloid and they remove thyroid hormones from thyroglobulin and them into circulation.

#### (b) IODINE ACCUMULATION BY THYROID :-

- ① Bio synthesis of iodinated hormones in the thyroid gland start with the entry of iodine ion from the blood. This entry <sup>take</sup> place against the concentration gradient, the follicle cells of the thyroid is able to trap iodine enormously higher amount (several hundred times) than that in circulation. The transport of iodine from the blood and its accumulation in the thyroid are accelerated by TSH while thiocyanate and perchlorate produce reverse effect.



(3)

② The thyroid follicular cells are able to trap iodide at the base of the cell and transport against an electrical gradient across the cell. Accumulation of iodine by the thyroid is an energy requiring mechanism. Its  $\text{Na}^+$  dependence was soon recognized and a  $\text{Na}^+ - \text{I}^-$  cotransport system was demonstrated. The  $\text{Na}^+ - \text{I}^-$  cotransport is inserted into the base lateral membrane of thyroid cell (thyrocytes) its activity is dependent on the  $\text{Na}^+$  gradient across the membrane and therefore on  $\text{Na}^+/\text{K}^+$ -ATPase. Therefore iodine is accumulated in gland. The gland contain iodide of 25 to 40 times its concentration in plasma and rapidly converts it to organically bound iodine within about 10 sec. Normally the thyroid contain approximately 10 mg of exchangeable iodine compared with 5000 - 7000 mg organically bound iodine.

④  
© OXIDATION OF IODIDE :-

The thyroid promptly undergoes oxidation to an active form probably the iodonium ion ( $I^+$ ) or Hypoiodite ( $HIO$ ) or both in the presence of  $H_2O_2$  and enzyme iodide peroxidase. The thyroidal peroxidase may be present within the golgi derived vesicle containing newly synthesized TG.

④ ➤ Formation of organically bound IODINE COMPOUND:-

Conversion of iodine to active iodide and organification of iodide into active tyrosine residue of Thyroglobulin may occurs as TG is secreted into the follicular lumen. The oxidized iodide is then incorporated into tyrosyl group of thyroglobulin the synthesis of Thyroxin in TG involves as -

1. The active iodide react with tyrosine molecule of TG incorporated of single active iodide into the phenolic ring of tyrosine yield 3 moniodotyrosine(MIT)
2. A second iodide may be incorporated into the 5<sup>th</sup> position to yield 3,5 diiodotyrosine (DIT).
3. Two molecule of the latter DIT undergo oxidative coupling to form tetra iodothyronine (Thyroxine) with the elimination of one alanine side chain.
4. Simuntaneously , one molecule each of Mono and di-iodotyrosines may couple to form triiodothyronine. Two molecule of moniodotyrosine form diiodothyronine.



(5)



These iodothyronines is peptide linkage with the globulin constitute the thyroglobulin molecule which is stored in the colloid material within the lumen of the thyroid follicles. The iodination and coupling reaction are accelerated by TSH and inhibited by certain anti-thyroid agents (e.g. thiocarbamides, amino benzenes etc.)

### e) PROTEOLYSIS OF THYROGLOBULIN :-



In accordance with demand, the thyroglobulin undergoes hydrolysis by a proteolytic enzyme (proteinases) to release the iodinated compound which then exhibit the hormonal activity. This process again accelerate by TSH.

Colloid is engulfed by follicular cell (Pseudopods) and transported into the cell as colloid droplets. These colloid containing vesicle then fuse with lysosome and are referred to as secondary lysosome. Much of the T<sub>4</sub> within these vesicle is degraded by lysosomal proteolytic enzymes. The thyroid hormones are then released (presumably by diffusion) into the cytoplasm and enter the extracellular space by diffusion, through the basal or lateral follicular membrane through Exocytosis of vesicular products, including T<sub>3</sub> and T<sub>4</sub> is not excluded. The iodinated tyrosine Tyrosomal proteolysis are then diiodinated by di-iodinase and recycled for are within the cell. Fig Please turn over

(6)



3,5-3'-Tetraiodothyronine (Thyroxine)  
 (contains 35-40% of thyroid iodine)



Fig

Biosynthesis of Thyroid hormone

